



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2012

---

## **The 'Effects of Transfusion Thresholds on Neurocognitive Outcome of Extremely Low Birth-Weight Infants (ETTNO)' Study: Background, aims, and study protocol**

Franz, A R ; Maier, R F ; Thome, U H ; et al ; Bucher, H U

**Abstract:** Background: Infants with extremely low birth weight uniformly develop anemia of prematurity and frequently require red blood cell transfusions (RBCTs). Although RBCT is widely practiced, the indications remain controversial in the absence of conclusive data on the long-term effects of RBCT. Objectives: To summarize the current equipoise and to outline the study protocol of the 'Effects of Transfusion Thresholds on Neurocognitive Outcome of extremely low birth-weight infants (ETTNO)' study. Methods: Review of the literature and design of a large pragmatic randomized controlled trial of restrictive versus liberal RBCT guidelines enrolling 920 infants with birth weights of 400-999 g with long-term neurodevelopmental follow-up. Results and Conclusions: The results of ETTNO will provide definite data about the efficacy and safety of restrictive versus liberal RBCT guidelines in very preterm infants.

DOI: <https://doi.org/10.1159/000335030>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-74216>

Journal Article

Published Version

Originally published at:

Franz, A R; Maier, R F; Thome, U H; et al; Bucher, H U (2012). The 'Effects of Transfusion Thresholds on Neurocognitive Outcome of Extremely Low Birth-Weight Infants (ETTNO)' Study: Background, aims, and study protocol. *Neonatology*, 101(4):301-305.

DOI: <https://doi.org/10.1159/000335030>

# The 'Effects of Transfusion Thresholds on Neurocognitive Outcome of Extremely Low Birth-Weight Infants (ETTNO)' Study: Background, Aims, and Study Protocol

ETTNO Investigators<sup>1</sup>

## Key Words

Infant, preterm • Anemia of prematurity, treatment • Transfusion • Randomized controlled trial

## Abstract

**Background:** Infants with extremely low birth weight uniformly develop anemia of prematurity and frequently require red blood cell transfusions (RBCTs). Although RBCT is widely practiced, the indications remain controversial in the absence of conclusive data on the long-term effects of RBCT. **Objectives:** To summarize the current equipoise and to outline the study protocol of the 'Effects of Transfusion Thresholds on Neurocognitive Outcome of extremely low birth-weight infants (ETTNO)' study. **Methods:** Review of the literature and design of a large pragmatic randomized controlled trial of restrictive versus liberal RBCT guidelines enrolling 920 infants with birth weights of 400–999 g with long-term neurodevelopmental follow-up. **Results and Conclusions:** The results of ETTNO will provide definite data about the efficacy and safety of restrictive versus liberal RBCT guidelines in very preterm infants.

Copyright © 2012 S. Karger AG, Basel

## Introduction

Infants with extremely low birth weight uniformly develop anemia of prematurity and frequently require multiple red blood cell transfusions (RBCTs) during neonatal intensive care [1–3]. The criteria currently applied to indicate RBCTs in this population are based on expert opinion rather than evidence and there is a lack of conclusive data on long-term effects of RBCT practices. Transfusing infants to improve oxygen-carrying capacity and restricting RBCT to avoid transfusion-associated risks and costs may both potentially impair long-term development.

## Potential Risks and Benefits of RBCTs

Although RBCTs are a universally accepted part of the treatment of preterm infants, recent reviews concluded that there are insufficient data on the long-term effects of RBCT practices [4, 5]. In the absence of uniformly accepted physiologic or evidence-based RBCT criteria, RBCT practices vary greatly between neonatal intensive-care units [1, 6–10].

RBCTs may be associated with retinopathy of prematurity [11–13], bronchopulmonary dysplasia [14, 15],

<sup>1</sup> The names and affiliations of the ETTNO Investigators are listed in the Appendix.

acute pulmonary deteriorations [16], necrotizing enterocolitis [17–19], intraventricular hemorrhage [20], and increased mortality [21] although these associations require cautious interpretation because of the retrospective and observational character of the underlying studies [22]. RBCTs are also associated with a minute residual risk of transmission of infectious agents [23]. Implementing RBCT guidelines effectively reduces the number of RBCTs administered to preterm infants [24, 25], and ‘restrictive’ RBCT guidelines are feasible in very low birth-weight infants and may result in very low numbers and low cumulative volumes of RBCTs [26].

However, reducing RBCTs by accepting low hemoglobin concentrations carries the risk of an at least temporarily insufficient oxygen transport to vital organs and impaired outcome [27].

In a recent randomized trial of restrictive versus liberal RBCT guidelines in 103 infants with birth weights of 500–1,300 g, restrictive RBCT guidelines were associated with a marginally increased incidence of apnea [28]. The study reported an increased incidence of brain injury (intraventricular hemorrhage grade 4 or periventricular leukomalacia) with restrictive RBCT guidelines. This study was criticized because (a) the primary endpoint for sample size determination was the number of transfusions and not a clinically relevant outcome parameter, (b) the combined outcome of intraventricular hemorrhage grade 4 or periventricular leukomalacia had not been a pre-defined outcome measure, (c) intraventricular hemorrhage usually occurs at a time when infants had not yet been enrolled into that study, (d) there was an excess of male patients in the restrictive-transfusion group and male infants tend to have poorer outcomes, (e) only 52% of the patients had a late cranial ultrasound, despite the latter being required to assess the true incidence of periventricular leukomalacia [29–31]. In contrast to the initial findings, long-term follow-up showed poorer results in measures of associative verbal fluency, visual memory, and reading in the liberally transfused group [32] and liberally transfused girls had the most prominent abnormalities on magnetic resonance imaging [33]. Again, these latter results have to be interpreted with caution because only about half of the patients initially enrolled in the trial completed these follow-up examinations.

In another recent randomized trial of restrictive versus liberal RBCT guidelines, the Premature Infant in Need of Transfusion (PINT) Trial, 451 infants with a birth weight <1,000 g were enrolled. Restrictive RBCT guidelines were associated with fewer transfusions but there were no differences in the primary outcome, death

or survival with retinopathy of prematurity  $\geq 3$ rd degree, bronchopulmonary dysplasia, or brain injury on ultrasound [34]. Nor was there a statistically significant difference in the incidence of death or severe adverse neurodevelopmental outcome [35]. However, post-hoc analyses showed that the proportion of infants with cognitive delay defined as a Mental Developmental Index (MDI) <85 was lower and the mean MDI was higher in the liberal-transfusion group, raising concern that restrictive transfusion guidelines may result in impaired neurodevelopment. Notably, maintaining higher hemoglobin concentrations appeared to be less costly [36].

This study was criticized for the fact that the mean difference in hemoglobin levels between the two treatment groups was only marginal (1 g/dl) and that the mean hemoglobin levels were high in both groups (10 vs. 11 g/dl) and may not reflect the range of RBCT guidelines currently proposed [37].

## Open Questions

There is insufficient evidence to decide whether preterm infants should be treated according to liberal or restrictive RBCT guidelines [4, 5]. The long-term safety and efficacy of ‘restrictive’ RBCT practices can only be evaluated in an adequately powered, large, randomized controlled trial with long-term neurodevelopmental follow-up and with a sufficient difference in mean hemoglobin levels between both treatment arms to reflect the range of RBCT guidelines currently applied. Such a trial may help to improve survival without neurodevelopmental impairment in preterm infants.

## Summary of Study

The Effects of Transfusion Thresholds on Neurocognitive Outcome of Extremely Low Birth-Weight Infants (ETTNO) study is an observer-blinded randomized controlled clinical trial. Nine hundred and twenty infants of 400–999 g birth weight will be randomized to restrictive or liberal transfusion trigger thresholds between 48 and 72 h of life, stratified by center and birth weight (400–749 g/750–999 g) (table 1). Sample size calculations were based on a  $\chi^2$  test assuming a power of 80%, a two-sided significance level of 5%, an incidence of death or major neurodevelopmental impairment (where cognitive delay is defined as MDI <85) of 61% versus 51% in the restrictive-threshold and the liberal-threshold group, respec-

**Table 1.** ETTNO transfusion thresholds

Time after randomization	Hematocrit, %			
	restrictive RBCT thresholds		liberal RBCT thresholds	
	critical	noncritical	critical	noncritical
3–7 days of age	<34	<28	<41	<35
8–21 days of age	<30	<24	<37	<31
>21 days of age	<27	<21	<34	<28

‘Critical’ refers to state of health and is defined as the presence of any of the following: (1) requirement of mechanical ventilation (any mode, excluding continuous positive airway pressure), (2) requirement of continuous positive airway pressure with  $\text{FiO}_2 > 0.25$  for  $> 12$  h per 24 h, (3) patent ductus arteriosus requiring therapy, (4) more than 6 apneas that require stimulation per 24 h, or more than 4 desaturations to  $\text{SpO}_2 < 60\%$  per 24 h despite methylxanthines and continuous positive airway pressure and (5) acute sepsis or acute necrotizing enterocolitis requiring inotropic or vasopressor support.

tively, based on the reported outcome data of the PINT study [35].

The assigned RBCT guidelines will be applied until discharge home from hospital. Both, the restrictive and the liberal RBCT guidelines to be compared in this trial reflect current clinical practice [5, 6, 8, 26, 38–41]. These RBCT guidelines will result in a clinically relevant difference in mean hemoglobin concentrations between both treatment groups of about 2 g/dl and will therefore improve recognition of any effect of transfusion trigger thresholds on neurocognitive outcome compared with the above referenced PINT trial [34, 35].

The primary outcome, death or neurodevelopmental impairment, will be determined at 24 months of age, corrected for prematurity. Funding for longer-term follow-up until 5.5 years of age will be sought. Neurodevelopmental impairment will be defined as any of the following: cognitive delay defined as MDI score of the *Bayley Scales of Infant Development* (2nd edition)  $< 85$ , cerebral palsy, or severe visual or hearing impairment. In the intention-to-treat population of all randomized patients in whom the primary outcome is ascertained, the primary outcome variable will be analyzed by logistic regression with factors treatment, center, and birth weight category at a two-sided significance level of 5%.

The study has been approved by the leading ethics committee at the University of Tübingen, Germany, and by the German Regulatory Authority (Paul-Ehrlich-Institut). Written informed consent will be required from parents or caretakers.

Recruitment began in August 2011 and units are invited to participate. The results of ETTNO will provide

definite data about the efficacy and safety of restrictive versus liberal RBCT guidelines in very preterm infants.

Collaborations will be sought with other groups to perform individual patient data meta-analyses to (a) verify the effect of different levels of transfusion trigger thresholds on neurodevelopmental outcome and (b) analyze the impact of transfusion triggers on less frequently observed complications of prematurity, such as severe retinopathy, severe chronic lung disease, necrotizing enterocolitis and others.

## Appendix

### *The ETTNO Investigators*

Principal Coordinating Investigator: Axel R. Franz, Tübingen, Germany.

Steering Committee: Rolf F. Maier, Marburg; Ulrich H. Thome, Leipzig; Mario Rüdiger, Dresden; Martina Kron, Ulm; Dirk Bassler, Christian F. Poets, and Ingeborg Krägeloh-Mann, Tübingen, Germany.

Local Investigators: Mark Schoberer, Sonja Trepels-Kottek, Aachen; Gernot Buheitel, Wilfried Schenk, Augsburg; Christoph Bühner, Florian Guthmann, Lars Garten, Dieter Hüseman, Christoph Czernik, Berlin (Charité); Rainer Rossi, Thomas Kühn, Elisabeth Eilers, Berlin (Neukölln); Norbert Teig, Almut Weitkämper, Bochum, Germany; Jes Reinholdt, Gitte Veiergang, Copenhagen, Denmark; Claudia Roll, Wolfgang Pielemeier, Datteln; Violeta Ojinaga, Nicole Braun, Mario Rüdiger, Dresden; Hans-Jörg Bittrich, Kathrin Roefke, Erfurt; Michael Schroth, Hans-Georg Topf, Erlangen; Anja Stein, Matthias Keller, Britta-Maria Hüning, Essen; Doris Fischer, Horst Buxmann, Frankfurt/Main; Matthias Heckmann, Dirk Faas, Gießen; Axel von der Wense, Peter Gudowius, Hamburg (Altona); Sarah Kabisch, Georg Hillebrand, Hamburg (Eppendorf);

Bettina Bohnhorst, Corinna Peter, Florian Urlichs, Hannover; Marc Hoppenz, Thomas Pabst, Köln (städt. Kliniken); Angela Kribs, Ruth Maria Klein, Christoph Hünseler, Bernhard Roth, Anne Vierzig, Frank Eifinger, André Oberthür, Kathrin Mehler, Köln (Universität); Ulrich H. Thome, Corinna Gebauer, Matthias Knüpfer, Leipzig; Rolf F. Maier, Michael Zemlin, Marburg; Larissa Mähnhardt, Claudia Franziska Nußbaum, Alexandra Schwepcke, München; Georg Rellensmann, Esther Rieger-Fackeldey, Münster; Jan-Holger Schiffmann, Jens Grombach, Stefan Schäfer, Nürnberg; Hugo Segerer, Annette Keller-Wackerbauer, Regensburg; Rainer Burghard, Mechthild Hubert, Siegen; Matthias Vochem, Ulrich Pohlmann, Patrick Neuberger, Martin Wagner, Stuttgart; Axel Franz, Dirk Bassler, Tübingen; Hans Fuchs, Manuel Schmid, Reinhard Hopfner, Ulm, Germany.

DMC: Hans Ulrich Bucher, Zürich, Switzerland; Heike Rabe, Brighton, UK; Josef Högel, Ulm; Data Management: Corinna Engel, Joachim Riethmüller, Center for Pediatric Clinical Studies, Tübingen, Germany.

## Acknowledgements

The ETTNO investigators gratefully acknowledge the generous support by the Deutsche Forschungsgemeinschaft (grant: DFG Fr 1455/6-1) and Masimo Corp., Irvine, Calif., USA, which generously provides patient monitors.

## Identification

EudraCT-No.: 2010-021576-28. ClinicalTrials.gov Identifier: NCT01393496.

## Disclosure Statement

All ETTNO Investigators confirmed that they have no conflict of interest to declare.

## References

- 1 Bifano EM, Curran TR: Minimizing donor blood exposure in the neonatal intensive care unit. Current trends and future prospects. *Clin Perinatol* 1995;22:657–669.
- 2 Widness JA, Seward VJ, Kromer JJ, Burmeister LF, Bell EF, Strauss RG: Changing patterns of red blood cell transfusion in very low birth weight infants. *J Pediatr* 1996;129:680–687.
- 3 Asch J, Wedgwood JF: Optimizing the approach to anemia in the preterm infant: is there a role for erythropoietin therapy? *J Perinatol* 1997;17:276–282.
- 4 Bassler D, Weitz M, Bialkowski A, Poets CF: Restrictive versus liberal red blood cell transfusion strategies for preterm infants: a systematic review of randomized controlled trials. *Curr Pediatr Rev* 2008;4:143–150.
- 5 Crowley M, Kirpalani H: A rational approach to red blood cell transfusion in the neonatal ICU. *Curr Opin Pediatr* 2010;22:151–157.
- 6 Linderkamp O, Zilow EP, Zilow G: The critical hemoglobin value in newborn infants, infants and children (in German). *Beitr Infusionsther* 1992;30:235–246.
- 7 Blanchette VS, Hume HA, Levy GJ, Luban NL, Strauss RG: Guidelines for auditing pediatric blood transfusion practices. *Am J Dis Child* 1991;145:787–796.
- 8 Maier RF, Metzke B, Obladen M: Low degree of regionalization and high transfusion rates in very low birthweight infants: a survey in Germany. *J Perinat Med* 1998;26:43–48.
- 9 Bednarek FJ, Weisberger S, Richardson DK, Frantz IDr, Shah B, Rubin LP: Variations in blood transfusions among newborn intensive care units. SNAP II Study Group. *J Pediatr* 1998;133:601–607.
- 10 Ringer SA, Richardson DK, Sacher RA, Kesler M, Churchill WH: Variations in transfusion practice in neonatal intensive care. *Pediatrics* 1998;101:194–200.
- 11 Cooke RW, Clark D, Hickey-Dwyer M, Weindling AM: The apparent role of blood transfusions in the development of retinopathy of prematurity. *Eur J Pediatr* 1993;152:833–836.
- 12 Hesse L, Eberl W, Schlaud M, Poets CF: Blood transfusion. Iron load and retinopathy of prematurity. *Eur J Pediatr* 1997;156:465–470.
- 13 Inder TE, Clemett RS, Austin NC, Graham P, Darlow BA: High iron status in very low birth weight infants is associated with an increased risk of retinopathy of prematurity. *J Pediatr* 1997;131:541–544.
- 14 Cooke RW, Drury JA, Yoxall CW, James C: Blood transfusion and chronic lung disease in preterm infants. *Eur J Pediatr* 1997;156:47–50.
- 15 Silvers KM, Gibson AT, Russell JM, Powers HJ: Antioxidant activity, packed cell transfusions, and outcome in premature infants. *Arch Dis Child Fetal Neonatal Ed* 1998;78:F214–F219.
- 16 Joshi VH, Joshi PV, Webert K, Cairnie J, Sabourin T, Blajchman M, Kirpalani H: Does transfusion related acute lung injury (TRALI) occur in newborns? PAS-Abstr 2006;59:460.
- 17 Paul DA, Mackley A, Novitsky A, Zhao Y, Brooks A, Locke RG: Increased odds of necrotizing enterocolitis after transfusion of red blood cells in premature infants. *Pediatrics* 2011;127:635–641.
- 18 Blau J, Calo JM, Dozor D, Sutton M, Alpan G, La Gamma EF: Transfusion-related acute gut injury: necrotizing enterocolitis in very low birth weight neonates after packed red blood cell transfusion. *J Pediatr* 2011;158:403–409.
- 19 Christensen RD, Lambert DK, Henry E, Wiedmeier SE, Snow GL, Baer VL, Gerday E, Ilstrup S, Pysher TJ: Is 'transfusion-associated necrotizing enterocolitis' an authentic pathogenic entity? *Transfusion* 2010;50:1106–1112.
- 20 Baer VL, Lambert DK, Henry E, Snow GL, Butler A, Christensen RD: Among very-low-birth-weight neonates is red blood cell transfusion an independent risk factor for subsequently developing a severe intraventricular hemorrhage? *Transfusion* 2011;51:1170–1178.
- 21 dos Santos AM, Guinsburg R, de Almeida MF, Procianny RS, Leone CR, Marba ST, Rugolo LM, Fiori HH, Lopes JM, Martinez FE: Red blood cell transfusions are independently associated with intra-hospital mortality in very low birth weight preterm infants. *J Pediatr* 2011;159:371–376 e1–3.
- 22 Kirpalani H, Whyte R: Truths, associations, and hypotheses. *J Pediatr* 2011;159:359–361.
- 23 Busch MP, Kleinman SH, Nemo GJ: Current and emerging infectious risks of blood transfusions. *JAMA* 2003;289:959–962.
- 24 Alagappan A, Shattuck KE, Malloy MH: Impact of transfusion guidelines on neonatal transfusions. *J Perinatol* 1998;18:92–97.
- 25 Baer VL, Henry E, Lambert DK, Stoddard RA, Wiedmeier SE, Eggert LD, Ilstrup S, Christensen RD: Implementing a program to improve compliance with neonatal intensive care unit transfusion guidelines was accompanied by a reduction in transfusion rate: a pre-post analysis within a multi-hospital health care system. *Transfusion* 2010;51:264–269.



- 26 Franz AR, Pohlandt F: Red blood cell transfusions in very and extremely low birth weight infants under restrictive transfusion guidelines: is exogenous erythropoietin necessary? *Arch Dis Child Fetal Neonatal Ed* 2001;84:F96–F100.
- 27 Strauss RG: Commentary: is it safe to limit allogeneic red blood cell transfusions to neonates? *Neonatology* 2008;93:217–222.
- 28 Bell EF, Strauss RG, Widness JA, Mahoney LT, Mock DM, Seward VJ, Cress GA, Johnson KJ, Kromer IJ, Zimmerman MB: Randomized trial of liberal versus restrictive guidelines for red blood cell transfusion in preterm infants. *Pediatrics* 2005;115:1685–1691.
- 29 Boedy RF, Mathew OP: Randomized trial of liberal versus restrictive guidelines for red blood cell transfusion in preterm infants. *Pediatrics* 2005;116:1048–1049.
- 30 Murray N, Roberts I, Stanworth S: Red blood cell transfusion in neonates. *Pediatrics* 2005;116:1609.
- 31 Swamy RS, Embleton ND: Red blood cell transfusions in preterm infants: is there a difference between restrictive and liberal criteria? *Pediatrics* 2006;117:257–258.
- 32 McCoy TE, Conrad AL, Richman LC, Lindgren SD, Nopoulos PC, Bell EF: Neurocognitive profiles of preterm infants randomly assigned to lower or higher hematocrit thresholds for transfusion. *Child Neuropsychol* 2011;1–21.
- 33 Nopoulos PC, Conrad AL, Bell EF, Strauss RG, Widness JA, Magnotta VA, Zimmerman MB, Georgieff MK, Lindgren SD, Richman LC: Long-term outcome of brain structure in premature infants: effects of liberal vs restricted red blood cell transfusions. *Arch Pediatr Adolesc Med* 2010;165:443–450.
- 34 Kirpalani H, Whyte RK, Andersen C, Asztalos EV, Heddle N, Blajchman MA, Peliowski A, Rios A, LaCorte M, Connelly R, Barrington K, Roberts RS: The Premature Infants in Need of Transfusion (PINT) study: a randomized, controlled trial of a restrictive (low) versus liberal (high) transfusion threshold for extremely low birth weight infants. *J Pediatr* 2006;149:301–307.
- 35 Whyte RK, Kirpalani H, Asztalos EV, Andersen C, Blajchman M, Heddle N, LaCorte M, Robertson CM, Clarke MC, Vincer MJ, Doyle LW, Roberts RS: Neurodevelopmental outcome of extremely low birth weight infants randomly assigned to restrictive or liberal hemoglobin thresholds for blood transfusion. *Pediatrics* 2009;123:207–213.
- 36 Kamholz KL, Dukhovny D, Kirpalani H, Whyte RK, Roberts RS, Wang N, Mao W, Zupancic JA: Economic evaluation alongside the Premature Infants in Need of Transfusion randomised controlled trial. *Arch Dis Child Fetal Neonatal Ed* 2011; epub ahead of print.
- 37 Bell EF: Transfusion thresholds for preterm infants: how low should we go? *J Pediatr* 2006;149:287–289.
- 38 Böhler T, Janecke A, Linderkamp O: Blood transfusion in late anemia of prematurity: effect on oxygen consumption, heart rate, and weight gain in otherwise healthy infants. *Infusionsther Transfusionsmed* 1994;21:376–379.
- 39 Maier RF, Sonntag J, Walka MM, Liu G, Metzke BC, Obladen M: Changing practices of red blood cell transfusions in infants with birth weights less than 1000 g. *J Pediatr* 2000;136:220–224.
- 40 Murray NA, Roberts IA: Neonatal transfusion practice. *Arch Dis Child Fetal Neonatal Ed* 2004;89:F101–F107.
- 41 Bell EF: When to transfuse preterm babies. *Arch Dis Child Fetal Neonatal Ed* 2008;93:F469–F473.